

MOTRIN® Professional Dosing Information for Infants & Children

Note: If possible, use weight to dose, otherwise use age. To arrive at the correct dose, weigh your child before giving MOTRIN®. A healthcare professional should be consulted for dosing in children under the age of six months.

Motrin[®] IB
Ibuprofen

Always ask your healthcare professional which product is right for your child.

		Your child's weight and age							
		0-5 mos	6-11 mos	12-23 mos	2-3 yrs	4-5 yrs	6-8 yrs	9-10 yrs	11 yrs
Age	Weight	6-11 lbs	12-17 lbs	18-23 lbs	24-35 lbs	36-47 lbs	48-59 lbs	60-71 lbs	72-95 lbs
 <p>Concentrated MOTRIN® Infants' Drops Berry & Berry dye-free flavors Active Ingredient: Ibuprofen 50mg (in each 1.25mL) Use only as directed.</p>		—	1.25mL	1.875mL	—	—	—	—	—
 <p>Children's MOTRIN® Suspension Berry, Grape, Tropical Punch, Bubblegum & Berry dye-free flavors Active Ingredient: Ibuprofen 100mg (in each 5mL or 1 tsp) Use only as directed.</p>		—	—	—	1 tsp or 5mL	1 1/2 tsp or 7.5mL	2 tsp or 10mL	2 1/2 tsp or 12.5mL	3 tsp or 15mL
 <p>MOTRIN® Junior Strength Easy-to-Swallow Caplets Active Ingredient: Ibuprofen 100mg (in each caplet) Use only as directed.</p>		—	—	—	—	—	2 caplets	2 1/2 caplets	3 caplets
 <p>MOTRIN® Junior Strength Chewable Caplets Orange & Grape flavors Active Ingredient: Ibuprofen 100mg (in each tablet) Use only as directed.</p>		—	—	—	1 tablet	1 1/2 tablets	2 tablets	2 1/2 tablets	3 tablets

tsp = teaspoon*

Take every 6-8 hours as needed. Do not exceed more than 4 doses in 24 hours.

Important Instructions for Proper Use

- Read and follow the label instructions on all MOTRIN® products
 - Take every 6-8 hours as needed
 - Do not exceed more than 4 doses in 24 hours
- Do NOT administer longer than 10 days, unless directed by a doctor
- Ask a doctor or pharmacist before use if the child is taking any other drug containing an NSAID (prescription or non-prescription)
 - MOTRIN® contains ibuprofen
- Keep all medicines out of the reach of children
- *Use only the dosing device that comes with a specific product
 - Household items such as measuring spoons are less accurate
- Do NOT administer adult medicines to children
- Concentrated MOTRIN® Infants' Drops are more concentrated than Children's MOTRIN® Liquids. The Concentrated Infants' Drops have been specifically designed for use with enclosed dosing device.
- Children's MOTRIN® Liquids are less concentrated than Concentrated MOTRIN® Infants' Drops. The Children's MOTRIN® Liquids have been specifically designed for use with the enclosed measuring cup.
- The OTC dosing: the recommended dose is 7.5mg/kg every 6-8 hours. The recommended daily dose is 30mg/kg.
- If you have any questions, contact your healthcare professional or call 1-877-895-3665